

23w

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
10 February 2005 (10.02.2005)

PCT

(10) International Publication Number  
**WO 2005/011813 A2**

(51) International Patent Classification<sup>7</sup>: **A61P 27/02**,  
37/06, 37/08, A61K 31/435, 31/436

(21) International Application Number:  
PCT/US2004/024054

(22) International Filing Date: 27 July 2004 (27.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
10/631,143 30 July 2003 (30.07.2003) US

(71) Applicant and

(72) Inventor: PEYMAN, Gholam, A. [US/US]; 8654  
Pontchartrain Blvd., Unit #1, New Orleans, LA 70124  
(US).

(74) Agents: LYMAN, Beverly, A. et al.; Wood, Herron &  
Evans, L.L.P., 2700 Carew Tower, Cincinnati, OH 45202  
(US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,  
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 2005/011813 A2

(54) Title: TREATMENT OF OCULAR DISEASE

(57) Abstract: A formulation to treat ocular conditions such as dry eye disease, as well as other conditions, is disclosed. Rapamycin and/or ascomycin is administered intraocularly, such as by topical application, injection into the eye, or implantation in or on the eye. For example, a topical administration may contain between about 50 pg/ml drug to about 50 µg/ml drug in a formulation which may be applied at bedtime or throughout the day. For injection, a dose of about 50 pg/ml to about 200 µg/ml may be used. Rapamycin and/or ascomycin may also be administered in milligram quantities as a surgical implant, for example, in a diffusible walled reservoir sutured to the wall of the sclera, or may be contained within an inert carrier such as microspheres or liposomes to provide a slow-release drug delivery system.

-1-

**TREATMENT OF OCULAR DISEASE****Field of the Invention**

The invention is directed to the use of rapamycin and ascomycin for therapeutic treatment of ocular conditions, including dry eye disease.

**Background**

5           Dry eye disease encompasses any condition where the tear film loses water and becomes more concentrated. It is a common complaint, affecting three million people in the United States alone, yet it is difficult to diagnose and treat. The loss of water from the tear film causes a corresponding rise in tear osmolarity. The increased osmolarity results in symptoms such as a sandy-gritty feeling in the eye, burning, irritation, or a foreign-body  
10   sensation that worsens during the day. Patients suffering from dry eye disease complain of mild to severe symptoms, with signs ranging from minimal superficial punctate keratitis to corneal perforation.

          Dry eye disease has a chronic remitting and relapsing nature and may result from a number of factors. The disease may be a natural part of the aging process, affecting  
15   15%-20% of adults over age 40. It may also result from pathological processes such as diseases of the lacrimal glands, mucus glands, and/or lipid producing glands, and may occur with cell infiltration or atrophy of the lacrimal gland (Sjögren's syndrome). Estrogen deficiency in postmenopausal women is also postulated to result in dry eye disease.

          One method to treat dry eye disease is by topical administration of over-the-  
20   counter drugs that serve as artificial tears. Numerous varieties of these artificial tears are

-2-

available (TheraTears® (Advanced Vision Research), Refresh® and Celluvisc® (Allergan), Tears Natural® and Bion Tears® (Alcon), GenTeal® and HypoTears® (CIBA Vision), each of which contain electrolytes and has varying pH levels, osmolarities, and surface tensions. Another method to treat dry eye disease is by surgery to close the lacrimal drainage ducts using punctum plugs. Neither method, however, is completely desirable. Artificial tears do not have a constant flow rate as do human tears, and treat the symptoms rather the cause of the disease. Surgery has its attendant risks, and may not be a viable option in older patients.

It is known that Cyclosporin A (cyclosporine, Allergan Inc.), may treat dry eye disease because patients administered Cyclosporin A for other disorders have shown a marked increase in tear flow. A topical formulation containing Cyclosporin A (Arrestase®, Allergan Inc.) is currently under review by the Food and Drug Administration. Cyclosporin A is an immunomodulator, suggesting that immune-mediated inflammation contributes to dry eye disease. Cyclosporin A has been used to treat various ocular pathologies such as glaucoma, corticosteroid-induced ocular hypertension, allograft rejection, infections, and ocular surface disease. It is also known that Cyclosporin A may be used in the eye to treat uveitis (inflammation of the uvea) by topical, intravitreal or systemic administration. Doses of 0.05%, 0.1%, and 0.5% Cyclosporin A have been reported. Cyclosporin A has good penetration into the cornea but not into the anterior chamber, and does not increase intraocular pressure or cause cataracts.

Tacrolimus (Prograf®, previously known as FK-506) is an immunomodulating drug that has been applied topically to treat a variety of dermatoses. Topical administration of tacrolimus at doses ranging from 0.03%-0.3% resulted in significant clinical improvement in atopic dermatitis after 2-3 weeks treatment, and tacrolimus treatment of other dermatologic diseases shows promise. Tacrolimus, like cyclosporine, blocks the signal transduction pathway needed to induce interleukin-2 gene expression and thereby activate T lymphocytes. In addition to suppressing T cell activation, tacrolimus inhibits anti-IgE-triggered histamine release and inhibits prostaglandin D2 synthesis in human skin mast cells. While oral administration produces limiting adverse effects (systemic immunosuppression, infection, neural toxicity, nephrotoxicity, and hypertension), topical administration for treatment of dermatoses at concentrations up to 0.3% showed no significant difference in effects between treated and

control groups. In addition, tacrolimus is well tolerated locally and only occasionally causes mild irritation. The use of tacrolimus to treat a variety of ocular conditions, including dry eye disease, has been reported in U.S. Patent No. 6,489,335.

Other compositions and methods to treat ocular conditions and enhance ocular lubrication would be desirable.

#### Summary of the Invention

A method to treat an ocular condition in a patient by intraocularly administering a pharmaceutically acceptable formulation of rapamycin in an amount effective to treat the condition. The method provides treatment while avoiding systemic administration of rapamycin.

10 In one embodiment, a sustained release pharmaceutically acceptable formulation is implanted intraocularly. For example, a matrix containing in the range of between about 3 mg rapamycin to about 5 mg rapamycin may be implanted in or on the eye and may continuously deliver rapamycin for ten or more years.

In another embodiment a concentration up to about 200 µg rapamycin is administered intraocularly without substantial toxicity. In another embodiment, rapamycin at a concentration in the range of about 1 ng/ml (0.0000001%) to less than 0.1 µg/ml (less than 0.001%) is administered topically. In other embodiments, rapamycin at a concentration in the range of about 1 ng/ml to about 200 µg/ml is injected under the conjunctiva, or a concentration in the range of about 1 ng/0.1 ml to about 200 µg/ml is injected in the vitreous, or a concentration in the range of about 20 µg/ml to about 200 µg/ml is injected behind the eyeball.

20 The method may prevent, decrease the time of onset, or lessen the severity of a wide variety of ocular conditions such as retinitis pigmentosa, ocular irritation following corneal surgery (e.g., LASIK surgery), age related macular degeneration, diabetic retinopathy, dry eye disease, scleritis, papillitis, and uveitis, as examples. Rapamycin may be administered in combination with Cyclosporin A and/or tacrolimus for intraocular administration without toxicity-limiting effects. Rapamycin may also be administered in combination with antibiotics without toxicity-limiting effects.

Another embodiment of the invention is a method to treat ocular conditions including ocular irritation following corneal surgery, retinitis pigmentosa, age related macular degeneration, diabetic retinopathy, scleritis, papillitis, and uveitis by intraocular administration of

ascomycin. Ascomycin may be the sole active agent or, alternatively, it may be administered in combination with other agents, including but not limited to rapamycin, Cyclosporin A, tacrolimus, and antibiotics.

In one embodiment, a sustained release pharmaceutically acceptable  
5 formulation is implanted intraocularly. For example, a matrix containing in the range of between about 3 mg ascomycin to about 5 mg ascomycin may be implanted in or on the eye and may continuously deliver ascomycin for ten or more years. In another embodiment a concentration up to about 200 µg ascomycin is administered intraocularly without substantial toxicity. In another embodiment, ascomycin at a concentration in the range of about 1 ng/ml (0.000001%)  
10 to less than 1 µg/ml (less than 0.0001%) is administered topically. In other embodiments, ascomycin at a concentration in the range of about 1 ng/ml to about 200 µg/ml is injected under the conjunctiva, or a concentration in the range of about 1 ng/0.1 ml to about 200 µg/ml is injected in the vitreous, or a concentration in the range of about 20 µg/ml to about 200 µg/ml is injected behind the eyeball.

15 The invention also encompasses a composition formulated for intraocular administration and dosing with rapamycin and/or ascomycin in a pharmaceutically acceptable formulation (e.g., in a physiologically acceptable solvent, buffered to a physiological pH, etc.). The composition may be in a solution, a suspension, an emulsion, etc., and it may be administered in the form of eye drops, a cream, an ointment, a gel, an injectable, etc., to the eye  
20 and/or the eye lid. The composition contains rapamycin and/or ascomycin in an amount effective to treat an ocular condition without substantial toxicity. As one example, the composition may contain rapamycin in a formulation for topical administration in an amount ranging from about 1 ng rapamycin to about 10 µg rapamycin. As another example, the composition may contain ascomycin in a formulation for topical administration containing be  
25 formulated as a cream containing ascomycin in an amount ranging from about 1 ng ascomycin to about 10 µg ascomycin. In one embodiment, a patient may self-administer the composition at a prescribed dosing interval (e.g., once a day, twice a day, as needed, etc.). In another embodiment, a medical professional may administer an intraocular injection. Alternatively, the composition may be a lipid or another type of matrix containing milligram quantities of

rapamycin and/or ascomycin. In this embodiment, the formulation is implanted in the eye and sustainedly releases the agent over an extended period of time.

These and other embodiments of the invention will be further appreciated with reference to the following detailed description.

## 5 Detailed Description

Methods are disclosed to ameliorate ocular conditions in need of treatment by intraocularly administering compositions containing rapamycin, ascomycin, and/or combinations of rapamycin and ascomycin as an active component. Intraocular administration provides direct effects to the eye while avoiding problems associated with systemic administration. Rapamycin is also known as sirolimus, RAPA and Rapamune. It is a triene macrolide antibiotic derived from *Streptomyces hydropiscus* and originally developed as an antifungal agent. Subsequently, it has shown anti-inflammatory, anti-tumor, and immunosuppressive properties. Ascomycin is also known as pimecrolimus, Immunomycin, and FR-900520. It is an ethyl analog of tacrolimus (FK506) and has strong immunosuppressant properties. It inhibits Th1 and Th2 cytokines, and preferentially inhibits activation of mast cells, and is used to treat contact dermatitis and other dermatological conditions. Rapamycin and ascomycin are commercially available, e.g., A.G. Scientific, Inc. (San Diego, CA).

Regarding its immunosuppressive potential, rapamycin has some synergetic effect with Cyclosporin A. It has been reported that rapamycin has a different mode of action compared to Cyclosporin A and tacrolimus, two other immunosuppressants. All three agents are immunosuppressants which affect the action of immune cell modulators (cytokines), but do not affect the immune cells themselves. However, while all three agents affect immune cell modulators, they do so differently: Cyclosporin A and tacrolimus prevent synthesis of cytokine messengers, specifically interleukin-2, while rapamycin acts on cytokine that has already been synthesized, preventing it from reaching immune cells.

Rapamycin inhibits inflammation by acting on both T-lymphocytes and dendritic cells. The latter are the first cells to recognize antigens. Rapamycin blocks the growth of dendritic cells and a number of other cells, such as tumors and endothelial cells, which are activated by the tumor cell releasing vascular endothelial growth factor (VEGF). VEGF is a central regulator of angiogenesis (formation of new blood vessels from pre-existing vessels) and

vasculogenesis (development of embryonic vasculature through an influence on endothelial cell differentiation and organization). Diseases that are characterized by abnormal angiogenesis and vasculogenesis, such as some cancers and some ocular diseases, may show abnormal production of VEGF. Thus, control of VEGF function may be one means to control or treat these diseases. Rapamycin has also been used in the prevention of smooth muscle hyperplasia after coronary stent surgery.

Methods which include administering rapamycin or ascomycin, either alone or in combination, may be useful for alleviating the effects of conditions that result in lack of moisture or wetness in the eye. Two examples of pathological conditions resulting in dry eye include pemphigus and Sjögren's syndrome, which affect the eye by either damaging the conjunctival cells responsible for maintaining the wetness of the cornea and the conjunctiva, or by damaging the lacrimal glands of the eye and/or the meibomian glands of the eye lid. Other examples of pathological conditions resulting in dry eye include hypolacrimation, alacrima, xerophthalmia, Stevens-Johnson syndrome, pemphigus, ocular pemphigoid, marginal blepharitis, diabetes, and/or post-corneal surgery (including but not limited to post-LASIK surgery). Examples of non-pathological conditions resulting in dry eye include the aging process, environmental factors (e.g., dry home and/or work environments), and extended use of visual display terminals (e.g., employment, recreation).

The inventive method is also useful for patients with inflammation and in which generation of an immune response is implicated in ocular diseases or diseases having an ocular component. Examples include diabetic retinopathy, age-related macular degeneration, retinitis pigmentosa, and uveitis (anterior, intermediate, or posterior). In addition, inflammatory responses accompanying bacterial, fungal, and viral disease also affect the eye, and are amenable to methods of therapy using rapamycin.

Rapamycin alone, ascomycin alone, or a combination of rapamycin and ascomycin, may be intraocularly administered by any route. The agent(s) may be administered topically to the eye or eye lid, for example, using drops, an ointment, a cream, a gel, a suspension, etc. The agent(s) may be formulated with excipients such as methylcellulose, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyvinyl pyrrolidone, neutral poly(meth)acrylate esters, and other viscosity-enhancing agents. The agent(s) may be injected

into the eye, for example, injection under the conjunctiva or tenon capsule, intravitreal injection, or retrobulbar injection. The agent(s) may be administered with a slow release drug delivery system, such as polymers, matrices, microcapsules, or other delivery systems formulated from, for example, glycolic acid, lactic acid, combinations of glycolic and lactic acid, liposomes, silicone, polyanhydride polyvinyl acetate alone or in combination with polyethylene glycol, etc. The delivery device can be implanted intraocularly, for example, implanted under the conjunctiva, implanted in the wall of the eye, sutured to the sclera, for long-term drug delivery.

Rapamycin alone, ascomycin alone, or a combination of rapamycin and ascomycin may be administered in a topical formulation for treatment of ocular conditions. In one embodiment, the agent(s) is formulated for topical administration to stimulate tear production by administering a composition containing rapamycin at a concentration in the range of about 50 pg/ml (0.000000005%) to about 50 µg/ml (0.005%), ascomycin at a concentration in the range of about 50 pg/ml to about 50 µg/ml, or a combination of rapamycin and ascomycin to achieve a total concentration of both agents of about 50 pg/ml to about 50 µg/ml. Within this range the agent(s) has wide safety and efficacy, permitting specific doses or administration protocols to be formulated for specific applications. For example, some patients may prefer once a day administration compared to administration more than once a day, so a higher concentration of agent(s) may be used for these patients.

In another embodiment, rapamycin in amounts ranging from about 1 ng to about 10 µg is contained in an aqueous-based cream excipient. In another embodiment, ascomycin in amounts ranging from about 1 ng to about 10 µg is contained in an aqueous-based cream excipient. In another embodiment, rapamycin and ascomycin are both present in an aqueous-based cream excipient in various proportions to achieve a total amount of combined agents of about 1 ng to about 10 µg. The drug(s) may be incorporated directly into the cream in solution, or may be contained in liposomes or microspheres either in solution or in an anhydrous form. The cream formulation is usually applied to the eye at bedtime, but it may be applied any time throughout the day if the cream does not cause blurred vision. In another embodiment, the agent(s) is formulated as a solution or suspension and is applied topically in the form of eye drops.



Rapamycin may also be administered by injection. Intraocular injection may be desirable or necessary, for example, for conditions in which topical administration is either not advised or is inadequate, for patients who have difficulty self-administering medications, etc. In one embodiment, the volume injected is less than 0.3 ml. In another embodiment, the volume injected is in the range of about 0.01 ml to about 0.3 ml. For intravitreal administration (injection into the vitreous), rapamycin concentrations in the range of about 1 ng/0.1 ml to about 200 µg/ml (0.02%) may be used without toxicity or adverse side effects. In a specific embodiment, a dose of about 50 µg/0.1 ml is administered. Injection may also be subconjunctival (into the subconjunctiva), or retrobulbar (behind the eyeball). For subconjunctival injection, a dose in the range of about 1 ng/ml to about 200 µg/ml may be used. For retrobulbar injection, a dose in the range of about 20 µg/ml to about 200 µg/ml may be used. Ascomycin may also be formulated as an injectable for intraocular injection. The same doses as used for intraocular injection of rapamycin are used for intraocular injection of ascomycin.

For long term delivery of rapamycin and ascomycin, either alone or in combination, and/or for sustained release, a matrix housing or containing the agent(s) may be implanted into the eye. For example, a reservoir containing in the range of about 3 mg to about 5 mg of agent(s) is estimated to be able to release about 1 µg agent(s) per day. At such a release rate, continuous, sustained dosing may occur over 1000 to 5000 days. If less than 1 µg of agent(s) per day is released, sustained dosing may last up to or more than a decade. In one embodiment, less than 50 µg/day of agent(s) is released from the matrix. In another embodiment, agent(s) is released from the matrix at a rate in the range of about 50 pg/day to about 50 µg/day. In another embodiment, agent(s) is released from the matrix at a rate in the range of about 1 µg/day to about 5 µg/day.

A surgically implanted intraocular device or matrix may be a reservoir container having a diffusible wall of polyvinyl alcohol or polyvinyl acetate and containing milligram quantities of rapamycin, ascomycin, or a combination of rapamycin and ascomycin may be implanted in the sclera. As another example, milligram quantities of agent(s) may be incorporated into a polymeric matrix having dimensions of about 2 mm by 4 mm, and made of a polymer such as polycaprolactone, poly(glycolic) acid, poly(lactic) acid, or a polyanhydride, or a lipid such as sebacic acid, and may be implanted on the sclera or in the eye. This is usually

accomplished with the patient receiving either a topical or local anesthetic and using a small (3-4 mm incision) made behind the cornea. The matrix, containing the agent(s), is then inserted through the incision and sutured to the sclera using 9-0 nylon.

Rapamycin alone, ascomycin alone, or a combination of rapamycin and ascomycin may also be contained within an inert matrix for either topical application or injection into the eye. As one example of an inert matrix, liposomes may be prepared from dipalmitoyl phosphatidylcholine (DPPC), preferably prepared from egg phosphatidylcholine (PC) since this lipid has a low heat transition. Liposomes are made using standard procedures as known to one skilled in the art. The agent(s), in amounts ranging from nanogram to microgram quantities, is added to a solution of egg PC, and the lipophilic drug binds to the liposome.

A time-release drug delivery system may be implanted intraocularly to result in sustained release of the active agent(s) over a period of time. The implantable formation may be in the form of a capsule of any of the polymers previously disclosed (e.g., polycaprolactone, poly(glycolic) acid, poly(lactic) acid, polyanhydride) or lipids that may be formulated as microspheres. As an illustrative example, rapamycin may be mixed with polyvinyl alcohol (PVA), the mixture then dried and coated with ethylene vinyl acetate, then cooled again with PVA. Rapamycin bound with liposomes may be applied topically, either in the form of drops or as an aqueous based cream, or may be injected intraocularly. In a formulation for topical application, the drug is slowly released over time as the liposome capsule degrades due to wear and tear from the eye surface. In a formulation for intraocular injection, the liposome capsule degrades due to cellular digestion. Both of these formulations provide advantages of a slow release drug delivery system, allowing the patient a constant exposure to the drug over time.

In a time-release formulation, the microsphere, capsule, liposome, etc. may contain a concentration of rapamycin that could be toxic if administered as a bolus dose. The time-release administration, however, is formulated so that the concentration released at any period of time does not exceed a toxic amount. This is accomplished, for example, through various formulations of the vehicle (coated or uncoated microsphere, coated or uncoated capsule, lipid or polymer components, unilamellar or multilamellar structure, and combinations of the above, etc.). Other variables may include the patient's pharmacokinetic-pharmacodynamic parameters (e.g., body mass, gender, plasma clearance rate, hepatic function, etc.). The formation and

-10-

loading of microspheres, microcapsules, liposomes, etc. and their ocular implantation are standard techniques known by one skilled in the art, for example, the use a ganciclovir sustained-release implant to treat cytomegalovirus retinitis, disclosed in Vitreoretinal Surgical Techniques, Peyman et al., Eds. (Martin Dunitz, London 2001, chapter 45); Handbook of  
5 Pharmaceutical Controlled Release Technology, Wise, Ed. (Marcel Dekker, New York 2000), the relevant sections of which are incorporated by reference herein in their entirety.

As another example, rapamycin alone, ascomycin alone, or a combination of rapamycin and ascomycin may be dissolved in an organic solvent such as DMSO or alcohol as previously described and containing a polyanhydride, poly(glycolic) acid, poly(lactic) acid, or  
10 polycaprolactone polymer.

Rapamycin and/or ascomycin, either alone or in combination with immunosuppressant agents such as Cyclosporin A, tacrolimus, etc., may be administered intraocularly and without substantial toxicity, to treat ocular conditions associated with an autoimmune response and/or involving an antigen/antibody reaction. Exemplary diseases  
15 include retinopathy in diabetic patients, macular degeneration, retinitis pigmentosa, inflammatory diseases of the eye such as Behçet's syndrome, toxoplasmosis, Birdshot choroidopathy, histoplasmosis, pars planitis, sarcoidosis, inflammatory diseases of the choroid of unknown etiology such as sympathetic ophthalmia, serpiginous choroidopathy, diffuse pigment epitheliopathy, Vogt-Koyanagi syndrome, polyarteritis nodosa, juvenile rheumatic  
20 arthritis, other conditions of the eye including uveitis (inflammation of the uvea), scleritis, (inflammation of the sclera), neuritis (inflammation of the optic nerve), or papillitis (inflammation of the optic nerve head) using the methods and formulations previously described. This may be achieved by one or a combination of factors, such as by slowing disease progression, lessening its severity, lengthening the time of onset, etc.

25 The toxicity of rapamycin, which had been reported with rapamycin administered systemically, was thought to limit its intraocular use. Ocular toxicity may manifest as a gross and/or histologic retinal and/or vitreous toxic reaction. Evidence of such a toxic reaction may include one or more of white vitreous bodies, white vitreous opacities, electroretinography abnormalities such as reduction in mean B-wave amplitude in both scotopic  
30 and photopic conditions, occlusion of the temporal retinal vessels, and fibrin deposits.

In one embodiment, rapamycin alone, ascomycin alone, or a combination of rapamycin and ascomycin, is administered in an amount or at a dose that does not result in substantial toxicity to the eye. As used herein, a lack of substantial toxicity encompasses both the absence of any manifestations of toxicity, as well as manifestations of toxicity which one skilled in the art would consider not sufficiently detrimental to decrease or cease treatment. As one example, fibrin deposits may be present indicating some toxicity, but less than substantial toxicity if their duration, number, etc., as determined by one skilled in the art, does not warrant that treatment be stopped. As another example, white vitreous bodies and fibrin bodies may be present indicating some toxicity, but less than substantial toxicity if their duration, number, etc., as determined by one skilled in the art, does not warrant that treatment be stopped.

Intraocular administration of a dose up to about 200 µg of either rapamycin or ascomycin, or a combination of rapamycin and ascomycin to achieve a final dose of up to about 200 µg, does not result in substantial toxicity. This dose is similar to the dose of 250 µg tacrolimus for administration without substantial toxicity, reported in co-pending patent application Serial No. 10/247,220, which is expressly incorporated by reference herein in its entirety. The intravenous solution form of rapamycin may be diluted to achieve the indicated concentration using 0.9% NaCl or 5% dextrose, or an organic solvent such as dimethylsulfoxide (DMSO) or alcohol, preferably a low molecular weight alcohol. Intraocular administration may be any of the routes and formulations previously described. For injection, either a solution, emulsion, suspension of a liquid, capsular formulation of microspheres or liposomes, etc. may be used.

To treat uveitis, rapamycin may be injected subconjunctivally at a dose in the range of about 1 ng/ml to about 200 µg/ml, or intravitreally at a dose of about 1 ng/0.1 ml to about 200 µg/ml. In one embodiment, the dose is about 50 µg/0.1 ml. To treat scleritis involving the anterior sclera, rapamycin may be administered topically. To treat scleritis involving the posterior sclera, rapamycin may be administered by retrobulbar injection at a dose in the range of about 20 µg/ml to about 200 µg/ml and dissolved in DMSO or a low concentration of alcohol. To treat neuritis or papillitis, rapamycin may be administered by retrobulbar injection at a dose in the range of about 20 µg/ml to about 200 µg/ml. Ascomycin

-12-

may be administered by the same routes for the given indications using the same doses as for rapamycin.

Diabetic retinopathy is a leading cause of blindness. Patients with diabetes mellitus have an absolute or relative lack of circulating insulin and, through a variety of factors, frequently present with vascular changes in the retina. These changes manifest in retinal microaneurysms, small hemorrhages, and exudates, and lead to the formation of scar tissue. New blood vessels may form around the optic disk (proliferative retinopathy). Over time, the cumulative results of such vascular effects lead to ocular pathologies which, ultimately, decrease vision in the diabetic patient. Thus, compositions and methods which reduce these vascular changes, or reduce their effects, improve the chances of a diabetic patient either maintaining vision, or at least slowing loss of vision.

Macular degeneration, also called age related macular degeneration (AMD) is a pathological condition that results in proliferation of new blood vessels in the subretinal area. While the presence of the new vessels themselves is not problematic, new vessels leak blood and other serous fluid which accumulate in surrounding spaces. It is this fluid accumulation that leads to visual impairment. For example, in the retina, both the large vessels and the capillaries normally have intact vessel walls. In the choroid, the large vessels normally have intact vessel walls, but the capillary walls or membranes contain fenestrations or openings. Any endogenous or exogenous fluid present in these capillaries, for example, blood, serous fluid, solubilized drug, etc. will leak outside the vessels and into the surrounding area. The accumulation of fluid can result in serous and hemorrhagic detachment of the retinal pigment epithelium and neurosensory retina, and can lead to loss of vision due to fibrous deform scarring. Patients with an early stage of AMD can be diagnosed by the presence in the eye of abnormal clumps of pigments, termed drusen, which are dead outer segments of photoreceptor cells under the retinal pigment epithelium. The presence of large, soft drusen in the eye indicates a pre-stage of exudative AMD, and places these patients at higher-than-average risk for developing neovascularizations, especially if one eye is already affected.

Retinitis pigmentosa is a general term that encompasses a disparate group of disorders of rods and cones, which are the sensory structures in the retina. While retinitis pigmentosa is a genetic disorder, and is not an inflammatory process, one manifestation of the

disease is the presence of irregular black deposits of clumped pigment in the peripheral retina. Thus, there is likely at least some immune component to retinitis pigmentosa which treatment with an immunosuppressant drug may help to alleviate.

A possible mechanism for the therapeutic efficacy of rapamycin and ascomycin in ocular disease involves their immunosuppressant activity. For example, diabetic patients treated with immunosuppressant drugs for reasons unrelated to vision develop less retinopathy over time than other diabetic patients. As another example, the drusen that is present in AMD constitutes a chronic inflammatory stimulus that becomes the target for encapsulation by a variety of inflammatory mediators, such as complement. Treatment with immunosuppressant drug may ameliorate this reaction. Immunosuppressant therapy results in decreased numbers of circulating immunocompetent cells such as lymphocytes. These cells otherwise have the potential to participate in an immune response, to lodge within the small capillaries and arterioles of the eye to form blockages and hence occlude blood flow, etc. In addition to lymphocytes, other hematopoietic cells may also be affected by immunotherapy, and include erythrocytes (red blood cells), megakaryocytes (precursors to platelets) and thrombocytes (platelets), and other leukocytes (white blood cells), such as monocytes and granulocytes. Local or *in situ* administration of immunosuppressant agents to the eye would be expected to decrease the number of these cells, resulting in reduction in the immune response, less blockage, increased blood flow, and increased patency of the ocular vessels.

Rapamycin, ascomycin, or a combination of the two in any of the previously described formulations, dosages, compositions, routes of administration, etc. may be employed. The active ingredient may be rapamycin or ascomycin alone. In one embodiment, rapamycin may be administered as described in combination with one or more known immunosuppressant agents, such as Cyclosporin A and/or tacrolimus. In another embodiment, rapamycin may be administered as described in combination with one or more antibiotics. In another embodiment, ascomycin may be administered as described in combination with one or more known immunosuppressant agents. In another embodiment, ascomycin may be administered as described in combination with one or more antibiotics. Such a combination is useful for treating ocular diseases having a microbial component, for example, conjunctivitis. Antibiotics which may be used are known to one skilled in the art (see, for example, Goodman and Gilman's The

-14-

Pharmacological Basis of Therapeutics, Eighth Edition, Gilman et al., Eds. (Pergamon Press, New York 1990, pages 1024-1033, which is incorporated by reference herein). Such exemplary antibiotics are not limiting, and the invention encompasses antibiotics presently known and unknown in combination with rapamycin and/or ascomycin for treatment of ocular disease.

- 5 Because rapamycin is administered locally to the eye (e.g., intraocular injection, topical ocular application), the undesirable effects brought about by administration of systemic immunosuppressant therapy (e.g., decreased peripheral blood leukocyte count, susceptibility to infections, hepatic and renal toxicity of the immunosuppressant agent itself, etc.) are absent.

- 10 It should be understood that the embodiments of the present invention shown and described in the specification are only preferred embodiments of the inventor who is skilled in the art and are not limiting in any way. Therefore, various changes, modifications or alterations to these embodiments may be made or resorted to without departing from the spirit of the invention and the scope of the following claims.

What is claimed is:

-15-

1. A method to treat an ocular condition in a patient comprising intraocularly implanting a composition comprising a sustained release matrix and a drug selected from the group consisting of rapamycin, ascomycin, and combinations thereof in an amount effective to treat the condition.
2. The method of claim 1 to treat a condition selected from the group consisting of a condition associated with an immunologic reaction, an ocular age related condition, a ocular degenerative condition, a condition associated with ocular moisture, a post-corneal surgery condition, and combinations thereof.
3. The method of claim 1 to treat a condition selected from the group consisting of dry eye disease, scleritis, neuritis, papillitis, uveitis, retinopathy, retinitis pigmentosa, macular degeneration, pemphigus, Sjögren's syndrome, Behçet's syndrome, toxoplasmosis, Birdshot choroidopathy, histoplasmosis, pars planitis, sarcoidosis, sympathetic ophthalmia, serpiginous choroidopathy, diffuse pigment epitheliopathy, Vogt-Koyanagi syndrome, polyarteritis nodosa, and juvenile rheumatic arthritis.
4. The method of claim 1 wherein the composition is implanted on the sclera.
5. The method of claim 1 wherein the matrix contains in the range of about 3 mg of the drug to about 5 mg of the drug.
6. A method to treat an ocular condition in a patient comprising intraocularly administering a composition comprising a drug selected from the group consisting of rapamycin, ascomycin, and combinations thereof, the drug at a concentration up to about 200 µg in a pharmaceutically acceptable formulation effective to treat the condition without substantial toxicity.



-16-

7. The method of claim 6 to treat a condition selected from the group consisting of a condition associated with an immunologic reaction, an ocular age related condition, a ocular degenerative condition, a condition associated with ocular moisture, a post-corneal surgery condition, and combinations thereof.
8. The method of claim 6 to treat a condition selected from the group consisting of dry eye disease, scleritis, neuritis, papillitis, uveitis, retinopathy, retinitis pigmentosa, macular degeneration, pemphigus, Sjögren's syndrome, Behçet's syndrome, toxoplasmosis, Birdshot choroidopathy, histoplasmosis, pars planitis, sarcoidosis, sympathetic ophthalmia, serpiginous choroidopathy, diffuse pigment epitheliopathy, Vogt-Koyanagi syndrome, polyarteritis nodosa, and juvenile rheumatic arthritis.
9. The method of claim 6 wherein the composition is administered by a method chosen from topical application, intraocular injection, and intraocular implantation.
10. The method of claim 6 wherein the composition further comprises Cyclosporin A, tacrolimus, and combinations thereof.
11. A method to treat an ocular condition in a patient comprising intraocularly administering a composition consisting essentially of rapamycin in a pharmaceutically acceptable formulation effective to treat the condition by a method selected from the group consisting of topical administration at a concentration of about 50 pg/ml to about 50 µg/ml, subconjunctival injection at a dose in the range of about 1 ng/ml to about 200 µg/ml, intravitreal injection at a dose in the range of about 1 ng/0.1 ml to about 200 µg/ml, or retrobulbar injection at a dose in the range of about 20 µg/ml to about 200 µg/ml.
12. The method of claim 11 wherein injection is intravitreal at a dose of about 50 µg/0.1 ml.

-17-

13. The method of claim 11 to treat a condition selected from the group consisting of a condition associated with an immunologic reaction, an ocular age related condition, a ocular degenerative condition, a condition associated with ocular moisture, a post-corneal surgery condition, and combinations thereof.
14. The method of claim 11 to treat a condition selected from the group consisting of dry eye disease, scleritis, neuritis, papillitis, uveitis, retinopathy, retinitis pigmentosa, macular degeneration, pemphigus, Sjögren's syndrome, Behçet's syndrome, toxoplasmosis, Birdshot choroidopathy, histoplasmosis, pars planitis, sarcoidosis, sympathetic ophthalmia, serpiginous choroidopathy, diffuse pigment epitheliopathy, Vogt-Koyanagi syndrome, polyarteritis nodosa, and juvenile rheumatic arthritis.
15. An ocular treatment method comprising intraocularly administering to a patient after corneal surgery a composition consisting essentially of rapamycin in a pharmaceutically acceptable formulation and in an amount effective to enhance post-surgical ocular moisture in the patient.
16. The method of claim 15 wherein the composition is administered by a method selected from the group consisting of topical administration and intraocular injection.
17. The method of claim 15 wherein the composition is administered by subconjunctival injection at a dose in the range of about 1 ng/ml to about 200 µg/ml, intravitreal injection at a dose in the range of about 1 ng/0.1 ml to about 200 µg/ml, or retrobulbar injection at a dose in the range of about 20 µg/ml to about 200 µg/ml.
18. The method of claim 15 wherein the composition is implanted intraocularly.
19. The method of claim 15 wherein is administered topically at a concentration in the range between about 50 pg/ml to about 50 µg/ml.

-18-

20. An ocular treatment method comprising intraocularly administering to a patient after corneal surgery a composition consisting essentially of ascomycin in a pharmaceutically acceptable formulation and in an amount effective to enhance post-surgical ocular moisture in the patient.
21. The method of claim 20 wherein the composition is administered by a method selected from the group consisting of topical administration and intraocular injection.
22. The method of claim 20 wherein the composition is administered by subconjunctival injection at a dose in the range of about 1 ng/ml to about 200 µg/ml, intravitreal injection at a dose in the range of about 1 ng/0.1 ml to about 200 µg/ml, or retrobulbar injection at a dose in the range of about 20 µg/ml to about 200 µg/ml.
23. The method of claim 20 wherein the composition is implanted intraocularly.
24. The method of claim 20 wherein is administered topically at a concentration in the range between about 50 pg/ml to about 50 µg/ml.
25. A method to treat an ocular condition in a patient comprising intraocularly administering to the patient a pharmaceutically acceptable formulation of a drug selected from the group consisting of rapamycin, ascomycin, and combinations thereof, in an amount up to about 200 µg effective to treat an ocular condition without substantial toxicity and at least one  
5 antibiotic.
26. The method of claim 25 wherein the composition is administered by a route selected from the group consisting of topical administration, intraocular injection, and intraocular implantation.

-19-

27. A therapeutic composition for treating an ocular condition consisting essentially of rapamycin in a physiologically acceptable intraocular formulation and at a dose up to about 200 µg effective to treat the ocular condition without substantial toxicity.
28. The composition of claim 27 formulated for topical administration.
29. The composition of claim 27 formulated as an injectable.
30. A therapeutic composition for treating an ocular condition consisting essentially of ascomycin in a physiologically acceptable intraocular formulation and at a dose up to about 200 µg effective to treat the ocular condition without substantial toxicity.
31. The composition of claim 30 formulated for topical administration.
32. The composition of claim 30 formulated as an injectable.
33. A therapeutic composition for treating an ocular condition comprising a physiologically acceptable matrix and a drug selected from the group consisting of rapamycin, ascomycin, and combinations thereof in an amount ranging between 3 mg and 5 mg for intraocular implantation.
34. The composition of claim 33 wherein the matrix comprises a substance selected from the group consisting of lipid, polyvinyl alcohol, polyvinyl acetate, polycaprolactone, poly(glycolic)acid, poly(lactic)acid, and combinations thereof.
35. The composition of claim 33 wherein the matrix sustainedly releases the drug.

36. The composition of claim 33 wherein the matrix releases the drug at a rate selected from the group consisting of less than about 50 µg/day, in a range between about 50 pg/day to about 50 µg/day, and in a range between about 1 µg/day to about 5 µg/day.

37. A method to treat an ocular condition in a patient comprising intraocularly administering a composition consisting essentially of ascomycin in a pharmaceutically acceptable formulation effective to treat the condition by a method selected from the group consisting of topical administration at a concentration of about 50 pg/ml to about 50 µg/ml, subconjunctival  
5 injection at a dose in the range of about 1 ng/ml to about 200 µg/ml, intravitreal injection at a dose in the range of about 1 ng/0.1 ml to about 200 µg/ml, or retrobulbar injection at a dose in the range of about 20 µg/ml to about 200 µg/ml.

38. The method of claim 37 wherein injection is intravitreal at a dose of about 50 µg/0.1 ml.

39. The method of claim 37 to treat a condition selected from the group consisting of a condition associated with an immunologic reaction, an ocular age related condition, a ocular degenerative condition, a condition associated with ocular moisture, a post-corneal surgery condition, and combinations thereof.

40. The method of claim 37 to treat a condition selected from the group consisting of dry eye disease, scleritis, neuritis, papillitis, uveitis, retinopathy, retinitis pigmentosa, macular degeneration, pemphigus, Sjögren's syndrome, Behçet's syndrome, toxoplasmosis, Birdshot choroidopathy, histoplasmosis, pars planitis, sarcoidosis, sympathetic ophthalmia, serpiginous  
5 choroidopathy, diffuse pigment epitheliopathy, Vogt-Koyanagi syndrome, polyarteritis nodosa, and juvenile rheumatic arthritis.

**THIS PAGE BLANK (USPTO)**